

Curriculum Vita

Personal Background:

Family name: **izadi**

First name: **sadegh**

Gender: **male**

Date of Birth: year: **1973** month: **10** day: **23**

Place of Birth: **Noorabad**

Nationality: **Iranian**

Marital status: **married**

Work Address:

Neurology department, Shiraz University of Medical Sciences, Nemazee square, Motahhari clinic, Shiraz

I.R of Iran, Tel: 0098 711 6121065, Fax: 0098 711 6121065,
izadisad@sums.ac.ir

E-mail:

Educational background:

Date	Location	Degree
1992-1999	Shiraz Uni.(Iran)	MD
2000-2004	Shiraz Uni.(Iran)	Neurology

Employment:

2008-present Associate Professor of neurology, neurology department, Shiraz University of Medical Health Services, Shiraz, Iran. WWW.sums.ac.ir

Publications:

Epidemiology of stroke in Shiraz, Iran

[B Daneshfard, S Izadi, A Shariat...](#) - Iranian journal of ..., 2015 - ncbi.nlm.nih.gov

Background: Stroke is the main cause of physical disability and the second leading cause of death worldwide. Two-thirds of all strokes occur in the developing countries. Despite being preventable, stroke is increasingly becoming a major health issue in these countries. The ...

[Cited by 34](#)[Related articles](#)[All 12 versions](#)

[\[HTML\] nih.gov](#)[Full View](#)

Significant increase in the prevalence of multiple sclerosis in iran in 2011

[S Izadi, A Nikseresht, M Sharifian...](#) - Iranian journal of ..., 2014 - ncbi.nlm.nih.gov

Multiple sclerosis (MS) is a chronic demyelinating disease of the central nervous system and usually affects young women. It is the second most common cause of neurological disability in young adults after trauma. 1, 2 Previously, Iran was considered a low prevalence area, but ...

[Cited by 55](#)[Related articles](#)[All 15 versions](#)

Correlation of clinical grading, physical tests and nerve conduction study in carpal tunnel syndrome

[S Izadi, B Kardeh, SSH Hooshiar...](#) - ... journal of pain, 2018 - degruyter.com

Background and aims Carpal tunnel syndrome (CTS) is a common debilitating condition. As the reliability of CTS-specific physical tests and its clinical grading remain a matter of debate, we determined the correlations between these assessments with nerve conduction study ...

[Cited by 9](#)[Related articles](#)[All 4 versions](#)

Estimated prevalence and incidence of multiple sclerosis in Iran

[M Etemadifar, S Izadi, A Nikseresht, M Sharifian...](#) - European ..., 2014 - karger.com

Background: Multiple Sclerosis (MS) is a chronic demyelinating disease of the central nervous system and its epidemiology has changed in recent years. Previously, Iran was considered to have a low MS prevalence. In this study, we aimed to update the prevalence ...

[Cited by 74](#)[Related articles](#)[All 4 versions](#)

[\[HTML\] nih.gov](#)[Full View](#)

Prevalence and incidence of multiple sclerosis in Fars province, Southern Iran

[S Izadi, AR Nikseresht, M Poursadeghfard...](#) - Iranian journal of ..., 2015 - ncbi.nlm.nih.gov

Background: Multiple sclerosis (MS) is a chronic inflammatory and demyelinating disease of the central nervous system, which is accompanied by significant disabilities. Although there are some published data regarding the epidemiological features of MS in other parts of Iran ...

[Cited by 27](#)[Related articles](#)[All 19 versions](#)

[\[HTML\] sciencedirect.com](#)

Reasons for uncontrolled seizures in adults; the impact of pseudointractability

[AA Asadi-Pooya, M Emami, N Ashjazadeh...](#) - Seizure, 2013 - Elsevier

Purpose We investigated the various possible reasons for uncontrolled seizures in patients 18 years of age and older to determine the impact of pseudointractability. We also tried to investigate the various forms of pseudointractability. Methods In this cross-sectional study ...

[Cited by 34](#)[Related articles](#)[All 7 versions](#)

[\[HTML\] sciencedirect.com](#)

Treatment of refractory generalized convulsive status epilepticus with enteral topiramate in resource limited settings

[AA Asadi-Pooya, MJ Jahromi, S Izadi, Y Emami](#) - Seizure, 2015 - Elsevier

Purpose To explore the feasibility, safety and efficacy of enterally administered topiramate (TPM) as an adjunctive treatment for adult patients with refractory generalized convulsive status epilepticus (RGCSE). Methods This prospective open-label non-randomized clinical ...

[Cited by 16](#)[Related articles](#)[All 6 versions](#)

[\[HTML\] nih.gov](#)

[\[HTML\] Large-vessel stenosis in the patients with ischemic stroke in Iran: Prevalence, pattern, and risk factors](#)

[A Borhani-Haghighi, M Emami, AS Vasaksi... - journal of vascular ..., 2015 - ncbi.nlm.nih.gov](#)

Background Large artery disease (LAD) is a common cause of stroke, but a little is known regarding its role in Iranian stroke patients. The current study investigates the prevalence and risk factors for cervicocephalic arterial stenosis in the patients with ischemic stroke using ...

[Cited by 13](#)[Related articles](#)[All 14 versions](#)

[\[HTML\] nih.gov](#)

[\[HTML\] The prevalence and factors contributing to hemodynamic depression in patients undergoing carotid angioplasty and stenting](#)

[AB Haghighi, S Kokabi, S Yousefi... - Journal of vascular ..., 2015 - ncbi.nlm.nih.gov](#)

About 170 patients (67% male, mean age: 71+9.8, 55.9% right side, 82.9% symptomatic) were recruited. Mean degree of stenosis was 79.4% in operated side and 40.7% in nonoperated side. Predilation, postdilation, or both were conducted in 18(10.5%), 141(83%), 11(6.5%) patients ...

[Cited by 12](#)[Related articles](#)[All 11 versions](#)

[\[PDF\] wiley.com](#)[Full View](#)

[\[PDF\] Epidermal neural crest stem cell transplantation as a promising therapeutic strategy for ischemic stroke](#)

[MS Salehi, S Pandamooz, A Safari... - CNS Neuroscience ..., 2020 - Wiley Online Library](#)

Introduction Cell-based therapy is considered as promising strategy to cure stroke. However, employing appropriate type of stem cell to fulfill many therapeutic needs of cerebral ischemia is still challenging. In this regard, the current study was designed to ...

[Cited by 1](#)[Related articles](#)[All 5 versions](#)

[\[PDF\] Efficacy of whole extract of licorice in neurological improvement of patients after acute ischemic stroke](#)

[P Ravanfar, G Namazi, M Atigh, S Zafarmand... - Journal of Herbal ..., 2016 - Elsevier](#)

Objective Licorice root has been reported to contain several neuroprotective compounds. In the present study we investigated its benefit in the treatment of acute ischemic stroke for which, treatment modalities are limited. Design Randomized double-blind placebo ...

[Cited by 6](#)[Related articles](#)[All 5 versions](#)

[\[PDF\] researchgate.net](#)

[\[CITATION\] Clinical manifestation and factors associated with hospital mortality rate among patients with subarachnoid hemorrhage](#)

[A Bolouki, S Izadi, HR SHAHRAKI - Headache, 2019](#)

[Related articles](#)[All 4 versions](#)

[\[HTML\] karger.com](#)

[\[HTML\] Demographic and technical risk factors of 30-day stroke, myocardial infarction, and/or death in standard-and high-risk patients who underwent carotid angioplasty and ...](#)

[AB Haghighi, S Yousefi, E Bahramali, S Kokabi... - Interventional ..., 2014 - karger.com](#)

Background: Carotid angioplasty and stenting (CAS) is an accepted treatment to prevent stroke in patients with carotid artery stenosis. The purpose of this study is to identify risk factors for major complications after CAS. Materials and Methods: This is a prospective study ...

[Cited by 6](#)[Related articles](#)[All 13 versions](#)

[\[HTML\] The effects of modified anti-inflammatory diet on fatigue, quality of life, and inflammatory biomarkers in relapsing-remitting multiple sclerosis patients: A randomized ...](#)

[Z Mousavi-Shirazi-Fard, Z Mazloom... - International Journal ..., 2020 - Taylor & Francis](#)

Background: The role of dietary interventions in improving the symptoms of Multiple Sclerosis (MS) has always been considered, but few studies have been conducted in this area. This study aimed to investigate the effects of modified anti-inflammatory diet on fatigue ...

[Related articles](#)[All 3 versions](#)

[\[HTML\] nih.gov](#)

[\[HTML\] Prevalence of intracranial artery stenosis in Iranian patients with acute ischemic stroke using transcranial Doppler ultrasonography](#)

[A Shariat, L Niknam, S Izadi... - Iranian journal of neurology, 2016 - ncbi.nlm.nih.gov](#)

Background: The aim of this study is to determine the frequency of intracranial artery stenosis in patients with acute ischemic stroke in Iran. Methods: A total of 169 patients with acute ischemic stroke were eligible to participate and were enrolled in this study from ...

[Cited by 4](#)[Related articles](#)[All 8 versions](#)

[\[HTML\] nih.gov](#)

[\[HTML\] Evaluation of transesophageal echocardiography in detecting cardiac sources of emboli in ischemic stroke patients](#)

[MA Toodeji, S Izadi, A Shariat... - Medical journal of the ..., 2015 - ncbi.nlm.nih.gov](#)

Background: Embolus is one of the causes of ischemic stroke that can be due to cardiac sources such as valvular heart diseases and atrial fibrillation and atheroma of the aorta. Transesophageal echocardiography (TEE) is superior in identifying potential cardiac ...

[Cited by 3](#)[Related articles](#)[All 12 versions](#)

[\[HTML\] nih.gov](#)

[\[HTML\] A case of chronic inflammatory demyelinating polyneuropathy presented with unilateral ptosis](#)

[S Izadi, S Karamimaghham... - Medical journal of the ..., 2014 - ncbi.nlm.nih.gov](#)

Abstract Chronic Inflammatory Demyelinating Polyneuropathy is an autoimmune disease with progressive and relapsing courses. The main clinical presentations are diffuse deep tendon hyporeflexia or areflexia and symmetric proximal-distal muscles weakness ...

[Cited by 5](#)[Related articles](#)[All 15 versions](#)

[\[PDF\] sums.ac.ir](#)

[\[PDF\] Circulating Levels of Interleukin-10 and-17 in Patients with Cerebral Sinovenous Thrombosis \(CSVT\) in Acute and Subacute Stages: A Prospective Case-Control ...](#)

[S Izadi, A Borhani-Haghighi, K Bastani... - Iranian Journal of ..., 2018 - iji.sums.ac.ir](#)

Background: Cerebral sinovenous thrombosis (CSVT) is a neurovascular disorder that occurs when a blood clot develops in a vein near the brain. Evaluating the subsequent changes in inflammatory cytokines can better reveal the underlying pathogenesis ...

[Cited by 1](#)[Related articles](#)[All 12 versions](#)

[\[PDF\] sbmu.ac.ir](#)[Full View](#)

[\[PDF\] A case of Guillain-Barre syndrome presented with bilateral pseudo-internuclear ophthalmoplegia](#)

[S Izadi, I Sabzgolini - International Clinical Neuroscience Journal, 2015 - journals.sbmu.ac.ir](#)

Guillain-Barre syndrome (GBS) or acute idiopathic polyradiculoneuritis is an acquired immune-mediated inflammatory and mainly demyelinating disorder of the peripheral nervous system. Cranial nerves are affected in over 50% of all cases, with the facial nerves ...

[Cited by 1](#)[Related articles](#)[All 4 versions](#)

[\[HTML\] gmj.ir](#)

[\[HTML\] Prevalence of Late-Onset Multiple Sclerosis in Fars Province, Southern Iran](#)

[S Izadi, M Sharifian, A Nikseresht, S Rafiee - Galen Medical Journal, 2014 - gmj.ir](#)

Background: Multiple Sclerosis (MS) is the most common demyelinating and inflammatory disease of the central nervous system especially in young adults, but in a subgroup of patients the first clinical symptoms present after the age of 50. This clinical presentation is defined as ...

[Cited by 1](#)[Related articles](#)[All 3 versions](#)

[\[HTML\] Fingolimod versus natalizumab in patients with relapsing remitting multiple sclerosis: a cost-effectiveness and cost-utility study in Iran](#)

[M Rezaee, S Izadi, K Keshavarz... - Journal of medical ..., 2019 - Taylor & Francis](#)

Aims: Multiple sclerosis (MS) is a chronic, autoimmune, and inflammatory disease. If the first-line medicines are not effective enough, specialists will prescribe second-line medicines, such as natalizumab and fingolimod. This study aimed to compare the cost-effectiveness ...

Related articlesAll 4 versions

Assessing the Prescriptions of the Neurologists in Shiraz, Iran.

[S Izadi, L Niknam...](#) - Journal of Isfahan ..., 2013 - [search.ebscohost.com](#)

Background: Assessment of prescriptions is a simple way to obtain valuable information about health system in a community. We conducted a descriptive-analytic study to assess the prescriptions of the neurologists in Shiraz, Iran, during the first 6 months of 2012 ...

Cited by 1Related articlesAll 2 versions

[HTML] [nih.gov](#)

[HTML] Neuroprotective effect of dimethyl fumarate in stroke: The role of nuclear factor erythroid 2-related factor 2

[A Safari, H Badeli-Sarkala, MR Namavar...](#) - Iranian Journal of ..., 2019 - [ncbi.nlm.nih.gov](#)

Background: There is evidence that supports the neuroprotective effects of dimethyl fumarate (DMF) in stroke. Nuclear factor erythroid 2-related factor 2 (Nrf2) has both anti-oxidant and anti-inflammatory mechanisms. We investigated the neuroprotective effects of ...

Related articlesAll 7 versions

[PDF] [sid.ir](#)

Immunological correlates of adult onset idiopathic generalized tonic-clonic epilepsy before and after sodium valproate treatment

[PGH YOUSEFI, S Izadi, A Ghaderi](#) - 2004 - [sid.ir](#)

Objective: To investigate possible immunological humoral correlates in newly diagnosed adult-onset generalized tonic-clonic epilepsy among Iranian patients before and after sodium valproate treatment. Patients and Methods: 72 adult patients with newly diagnosed ...

Cited by 2Related articlesAll 15 versions

[PDF] [sbmu.ac.ir](#) Full View

A case of Guillain-Barre syndrome with significant lymphocytic pleocytosis

[S Izadi, M Karimian](#) - International Clinical Neuroscience Journal, 2017 - [journals.sbmu.ac.ir](#)

Guillain-Barre syndrome (GBS) is an immune-mediated polyneuropathy. In the cerebrospinal fluid (CSF) characteristic increased protein without pleocytosis ie, albuminocytologic dissociation is seen. But cases of GBS with significant CSF lymphocytosis ...

Cited by 1Related articlesAll 3 versions

The effect of self-care education on the quality of life of patient referring to Institute MS FarsIran, 2012

[F Ghodsbin, EP Manesh, Z Molazem...](#) - Advances in ..., 2013 - [go.gale.com](#)

Multiple sclerosis (MS) is a chronic and progressive disease of the central nervous system with debilitating symptoms and complications. Therefore, education and appropriate interventions, including self-care education, are necessary to increase the quality of life in ...

Cited by 1Related articles

[PDF] [gums.ac.ir](#)

Predictors and Conversion Rate of Clinically Isolated Syndrome to Clinically Definite Multiple Sclerosis: A Follow-up Study in Patients Living in the Southern Part of ...

[S Izadi, M Ahmadi, M Poursadeghfard](#) - Caspian Journal of ..., 2020 - [cjns.gums.ac.ir](#)

Background: Clinical course of Clinically Isolated Syndrome (CIS) is variable, and identifying patients who will eventually develop Multiple Sclerosis (MS) is essential. Objectives: To assess the conversion rate of CIS to Clinically Definite Multiple Sclerosis (CDMS) and its ...

[HTML] [emedicalj.com](#)

[HTML] Cost-effectiveness and Cost-utility of CinnoVex versus ReciGen in Patients with Relapsing-Remitting Multiple Sclerosis in Iran

[R Ravangard, M Rezaee, K Keshavarz...](#) - Shiraz E-Medical ..., 2018 - [emedicalj.com](#)

Background: Multiple Sclerosis (MS) is a chronic and inflammatory disease that can affect the patients' quality of life and impose many costs on them. Several types of medicine are used to change the course of the disease, treat disease-related attacks, and treat the ...

Related articlesAll 3 versions

Is there any relation between cervical cord plaques and discopathy in patients with multiple sclerosis?

[A Nikseresht](#), [M Sharifian](#), [S Izadi](#), [AH Jahromi](#)... - *Clinical Neurology and ...*, 2014 - Elsevier

Introduction Multiple sclerosis (MS) is the most common chronic autoimmune demyelinating disease of the central nervous system. The purpose of this study is to determine the relationship between the site of the cervical discopathy and cervical spinal cord plaque in ...

Cited by 1Related articlesAll 5 versions

[PDF] [sbmu.ac.ir](#)Full View

Evaluation of Serum Auto Antibodies in Multiple Sclerosis Patients: A Case Control Study

[S Izadi](#), [S Khoshniat](#) - *International Clinical Neuroscience ...*, 2017 - [journals.sbmu.ac.ir](#)

Background and Purpose: Multiple sclerosis (MS) is believed to be an immune-mediated disorder that develops from an interaction of the individual's genetic and as yet unidentified environmental causes. The prevalence of auto-antibodies in multiple sclerosis patients and ...

Related articlesAll 3 versions

[CITATION] *Journal of Herbal Medicine*

[P Ravanfar](#), [G Namazi](#), [M Atigh](#), [S Zafarmand](#)...

Related articles

[PDF] [researchgate.net](#)

[CITATION] Extracranial Carotid Artery Stenosis in patients with Ischemic Stroke: a hospital-based study in **Shiraz**, Iran

[Z BEIZAVI](#), [S IZADI](#), [B DANESHFARD](#), [A SHARIAT](#)

Related articlesAll 2 versions

Stroke Epidemiology in Southwestern Iran: a Hospital-Based Study in **Shiraz**

[B Daneshfard](#), [S Izadi](#), [A Shariat](#)... - *6th national & first ...*, 2013 - [researchgate.net](#)

Title: Stroke Epidemiology in Southwestern Iran: a Hospital-Based Study in Shiraz Authors:

Babak Daneshfard1, 2, Sadegh Izadi2, 3, Abdolhamid Shariat2, 3, Mohammad Amin

Toudaj1, 2, Zahra Beyzavi1, 2, Leila Niknam1, 2 Abstract: Background: Stroke is the second ...

All 2 versions

[HTML] [uswr.ac.ir](#)

[HTML] One year survival and quality of life in patients successfully discharged from neuro critical care unit

[M Poursadeghfard](#), [K Bastani](#)... - *Iranian Rehabilitation ...*, 2017 - [irj.uswr.ac.ir](#)

2. Methods This is a retrospective cohort study in a limited time. We enrolled all patients admitted to NICU in Nemazee Hospital, Shiraz, south of Iran (an educational center) over a period of one year (February 2011-February 2012). Nemazee Hospital has multiple ...

Related articlesAll 14 versions

[PDF] [iums.ac.ir](#)

[CITATION] CT-guided laser probe for ablation of brain tumors

[A Daneshi](#), [M Taghipour](#), [S Izadi](#)... - *Basic and ...*, 2010 - ... , Iran **University** of Medical Sciences

Related articlesAll 9 versions

[PDF] [karger.com](#)

[CITATION] Contents Vol. 3, 2014

[SA Quadri](#), [V Ramakrishnan](#), [O Hariri](#), [MA Taqi](#)... - *Interventional ...*, 2014 - [karger.com](#)

... Michael G. Hennerici Department of Neurology **University** of Heidelberg Mannheim, Germany ...

of 30-Day Stroke, Myocardial Infarction, and/or Death in Standard- and High-Risk Patients Who

Underwent Carotid Angioplasty and Stenting Borhani Haghghi, A. (**Shiraz**); Yousefi, S ...

[PDF] [academia.edu](#)

[\[PDF\] Predictors and Conversion Rate of Clinically Isolated Syndrome to Clinically Definite Multiple Sclerosis: A Follow-up Study in Patients Living in the Southern ...](#)

[S Izadi, M Ahmadi, M Poursadeghfard - 2020 - academia.edu](#)

Background: Clinical course of Clinically Isolated Syndrome (CIS) is variable, and identifying patients who will eventually develop Multiple Sclerosis (MS) is essential. Objectives: To assess the conversion rate of CIS to Clinically Definite Multiple Sclerosis (CDMS) and its ...

All 3 versions

[\[PDF\] pdfarchive.ir](#)

[\[PDF\] üvûÍ© û%o© xwÆΩν fl†¬ Ÿ ÆòÑŸ¤ wœ©†~ ñ¶› È• ùûz](#)

[È Áóí - pdfarchive.ir](#)

... Sciences, **Shiraz**, Iran 2- Resident, Department of Neurology, School of Medicine, **Shiraz University** of Medical Sciences, **Shiraz**, Iran 3- General Practitioner, School of Medicine, **Shiraz University** of Medical Sciences, **Shiraz**, Iran Corresponding Author: **Sadegh Izadi** MD, Email ...

Related articlesAll 4 versions

[\[PDF\] bpums.ac.ir](#)

[The Prevalence and Factors Contributing to Hemodynamic Depression in Patients Undergoing Carotid Angioplasty and Stenting](#)

[R Nemati, A Borhani Haghighi, S Kokabi - Journal of Vascular and ... - eprints.bpums.ac.ir](#)

... Shariat 1,2 , Alireza Nikseresht 1,2 , Nahid Ashjazadeh 1,2 , **Sadegh Izadi** 1,2 ... cross-sectional study conducted in Kowsar and Alzahra hospitals affiliated with **Shiraz Uni-** versity of ... study protocol was approved by the institutional review board (IRB) of **Shiraz University** of Medical ...

Congress Presentations:

1. **Sadegh Izadi**(1), Alireza Nikseresht(1), Maryam Sharifian .Significant increase in the prevalence of multiple sclerosis in Iran, a report on the patients receiving Beta Interferons in different provinces. **ECTRIMS 2012.lyon –france.**
2. **Izadi S.** Nikseresht AR, **Poursadeghfard M**, Yasdanparast R. epidemiology of multiple sclerosis in the cities of Fars province.19th congress of neurology. . 2012, May, Tehran, Iran.
3. **sadegh izadi.** A case of childhood traumatic epilepsy whit fecal incontinency. 8th international congress on epilepsy. 2011 November, Tehran, Iran.
4. **sadegh izadi .**A CASE OF CIDP WITH FLACTUATING PTOSIS. 17th international congress of neurology. 2010, Tehran, Iran.
5. **sadegh izadi .**Who much is the effective dose of antiepileptic drugs? 5th international congress on epilepsy. 2008, Tehran, Iran.
6. **sadegh izadi .**Comparison of prednisolon with acyclovir and prednisolon in treatment of Bell's palsy. 14th congress of neurology. . 2007, May, Tehran, Iran.

7. Sadegh izadi¹, Abas jaberi rahimi¹, Aliakbar Kadivar², K Kashani². A case of Concomitant central and peripheral nervous system demyelination disease. 16th congress of neurology. . 2009, May, Tehran, Iran.

8. Izadi S.1 *, MD, Nikseresht A.R 1, MD, kadivar A.1, MD. Epidemiology of Multiple Sclerosis in Fars Province, Southern Iran.

9. Sadegh izadi. Evaluation of neuromuscular disorders in patients of Behzisty centers. 16th congress of neurology. . 2009, May, Tehran, Iran.

Research interesting:

1. Multiple sclerosis
2. stroke